

# Polytope Approximation and the Mahler Volume

**Sunil Arya**

Hong Kong University of Science and Technology

**Guilherme D. da Fonseca**


Universidade Federal do Estado do Rio de Janeiro (UniRio)

**David M. Mount**

University of Maryland, College Park

SODA 2012, Kyoto, Japan

# The Mahler Volume



- $K$ : convex body
- **Polar body of  $K$** : set of points  $p$  such that  $p \cdot q \leq 1$  for  $q \in K$
- **Mahler volume of  $K$** : product of the volume of  $K$  and the volume of  $\text{polar}(K)$

Important for us:


The Mahler volume of  $K$  is bounded below by a constant [Kup08]

- A regular simplex attains the minimum volume [KR11]
- Vast literature for centrally symmetric convex bodies

# Polytope Approximation

## Problem description:

- **Input:** convex body  $K$  in  $d$ -dimensional space and parameter  $\varepsilon$
  - **Output:** polytope  $P$  which  $\varepsilon$ -approximates  $K$  with a small number of facets (alternatively, vertices)
- 
- Focus on Hausdorff metric in Euclidean spaces of constant dimension  $d$
  - Assume (without loss of generality) that  $\text{diam}(K) = 1$
  - Assume the width of  $K$  is at least  $\varepsilon$ , for otherwise the problem instance should be solved in a lower dimensional space.



# Uniform vs. Nonuniform Bounds

- Several algorithms to find the “best” polytope for a given input
- **How good** is this best polytope?

## Nonuniform bounds:


- Hold for  $\varepsilon \leq \varepsilon_0$ , where  $\varepsilon_0$  **depends on the input**
- Example: Gruber [Gru93] bounds the complexity  $n$  using the Gaussian curvature  $\kappa$  of the input

$$n = 1/\varepsilon^{(d-1)/2} \int_{\partial K} \sqrt{\kappa(x)} dx$$

## Uniform bounds:

- Hold for  $\varepsilon \leq \varepsilon_0$ , where  $\varepsilon_0$  is a **constant**
- Example: Dudley [Dud74] and Bronshteyn and Ivanov [BI76] bound the maximum number of facets/vertices as a function of  $\varepsilon$ ,  $d$ , and the diameter of the input

# Dudley's Approximation




Dudley, 1974:

A convex body  $K$  of diameter 1 can be  $\varepsilon$ -approximated by a polytope  $P$  with  $O(1/\varepsilon^{(d-1)/2})$  facets.

- Dudley's approximation is the best possible for **balls**
- It oversamples areas of very high and very low curvatures
- Intuition: A **skinny** body should be **easier** to approximate

# Dudley's Approximation




Dudley, 1974:

A convex body  $K$  of diameter 1 can be  $\epsilon$ -approximated by a polytope  $P$  with  $O(1/\epsilon^{(d-1)/2})$  facets.

- Dudley's approximation is the best possible for **balls**
- It oversamples areas of very high and very low curvatures
- Intuition: A **skinny** body should be **easier** to approximate

# Dudley's Approximation



Dudley, 1974:

A convex body  $K$  of diameter 1 can be  $\epsilon$ -approximated by a polytope  $P$  with  $O(1/\epsilon^{(d-1)/2})$  facets.

- Dudley's approximation is the best possible for **balls**
- It oversamples areas of very high and very low curvatures
- Intuition: A **skinny** body should be **easier** to approximate

# Our Result: Improved Polytope Approximation

Better uniform bound for *skinny* bodies:

A convex body  $K$  can be  $\varepsilon$ -approximated by a polytope  $P$  with  $\tilde{O}(\sqrt{\text{area}(K)}/\varepsilon^{(d-1)/2})$  facets (alternatively, vertices).

- Uses **area** instead of diameter
- Matches Dudley's bound up to a **log** factor when the body is fat
- Significant improvement for skinny bodies
- Analysis uses several new techniques for the problem (polarity, Mahler volume,  $\varepsilon$ -nets...)




# Impact to Other Problems

## Approximate polytope membership


- For the **same storage**, the query time is reduced to the **square root**!
- $\tilde{O}(1/\varepsilon^{(d-1)/8})$  query time with  $O(1/\varepsilon^{(d-1)/2})$  (Dudley's) storage

## Approximate nearest neighbor (ANN)

- ANN reduces to polytope membership [AFM11]
- Improved query time for storage between  $O(n/\varepsilon^{d/4})$  and  $O(n/\varepsilon^{d-1})$



# Dual Caps and $\varepsilon$ -Dual Caps




- **Dual cap**: portion of the boundary of  $K$  visible from a point  $q$
- **Width**: distance between  $K$  and  $q$
- **$\varepsilon$ -dual cap**: dual cap of width  $\varepsilon$
- A set  $N$  of points **stabs** all  $\varepsilon$ -dual caps if for every dual cap  $D$  we have  $D \cap N \neq \emptyset$

## Lemma:

If a set  $N$  of points stabs all  $\varepsilon$ -dual caps, then the polytope defined by tangent hyperplanes constructed at the points of  $N$  is an  $\varepsilon$ -approximation to  $K$ .

# Voronoi Patches



- The **Voronoi region**  $\text{Vor}(D)$  of a dual cap  $D$  is the set of points closer to  $D$  than to any other points of  $K$
- **Dudley sphere**: Sphere  $S$  of radius 3 centered at the origin
- The **Voronoi patch** of a dual cap  $D$  is the intersection  $\text{Vor}(D) \cap S$

# Mahler Comes In

We show that:


An  $\varepsilon$ -dual cap  $D$  and its Voronoi patch are related in a manner that is similar to the **polar** transform (up to an  $\varepsilon$ -scaling).

Using the fact that the **Mahler volume** is at least a constant:

**Key lemma:**

For any  $\varepsilon$ -dual cap  $D$ , the product of  $\text{area}(D)$  and  $\text{area}(\text{Vor}(D) \cap S)$  is  $\Omega(\varepsilon^{d-1})$ .

Less formally: If  $D$  has small area, then its Voronoi patch is large



# Stabbing Large $\varepsilon$ -Dual Caps

- Large  $\varepsilon$ -dual cap:

$$\text{area}(D) \geq \sqrt{\text{area}(K)} \cdot \varepsilon^{(d-1)/2}$$

- Fraction of the boundary of  $K$  covered by  $D$ :

$$\alpha = \frac{\sqrt{\text{area}(K)} \cdot \varepsilon^{(d-1)/2}}{\text{area}(K)} = \frac{\varepsilon^{(d-1)/2}}{\sqrt{\text{area}(K)}}$$

- We stab large  $\varepsilon$ -dual caps with an  $\alpha$ -net on the boundary of the **convex body**
- Using VC-dimension arguments the size of the  $\alpha$ -net is

$$O\left(\frac{1}{\alpha} \log \frac{1}{\alpha}\right) = \tilde{O}\left(\frac{\sqrt{\text{area}(K)}}{\varepsilon^{(d-1)/2}}\right)$$

# Stabbing Small $\varepsilon$ -Dual Caps

- **Small**  $\varepsilon$ -dual cap:  $\text{area}(D) < \sqrt{\text{area}(K)} \cdot \varepsilon^{(d-1)/2}$
- By the key lemma, the Voronoi patch of  $D$  is large:

$$\text{area}(\text{Vor}(D) \cap S) > \frac{\varepsilon^{(d-1)/2}}{\sqrt{\text{area}(K)}}$$

- Dudley ball  $S$  has constant area
- Fraction of the boundary of  $S$  covered by  $\text{Vor}(D) \cap S$ :

$$\alpha = \frac{\varepsilon^{(d-1)/2}}{\sqrt{\text{area}(K)}}$$

- We stab small  $\varepsilon$ -dual caps indirectly, using an  $\alpha$ -net on the boundary of the **Dudley ball** and mapping the points back to  $K$
- Using VC-dimension arguments the size of the  $\alpha$ -net is

$$O\left(\frac{1}{\alpha} \log \frac{1}{\alpha}\right) = \tilde{O}\left(\frac{\sqrt{\text{area}(K)}}{\varepsilon^{(d-1)/2}}\right)$$

# Approximate Polytope Membership Queries

The **same methods** improve the existing bounds [AFM11] for:


## Polytope membership queries

Given a polytope  $P$  in  $d$ -dimensional space, **preprocess**  $P$  to answer **membership queries**:

Given a point  $q$ , is  $q \in P$ ?

## Approximate version

- An **approximation parameter**  $\epsilon$  is given (at preprocessing time)
- Assume the polytope has **diameter 1**
- If the query point's distance from  $P$ 's boundary:
  - $> \epsilon$ : answer must be **correct**
  - $\leq \epsilon$ : **either** answer is acceptable



# Approximate Polytope Membership Queries

Improved tradeoff for approximate polytope membership queries


**New bounds:**

For integer  $k \geq 2$ , we can answer  $\varepsilon$ -approximate polytope membership queries with

**Storage:**  $O(1/\varepsilon^{(d-1)/(1-k/2^k)})$

**Query time:**  $O(1/\varepsilon^{(d-1)/2^{k+1}} \log(1/\varepsilon))$

- For the same storage, the query time is reduced to roughly the **square root**
- Leads to improved approximate nearest neighbor data structures




**Storage:**  $O(1/\varepsilon^{(d-1)/2})$

**Query time:**  $\tilde{O}(1/\varepsilon^{(d-1)/8})$



# Approximate Nearest Neighbor (ANN) Searching



- **ANN**: Preprocess  $n$  points such that, given a query point  $q$ , can find a point within at most  $1 + \epsilon$  times the distance to  $q$ 's nearest neighbor
- It is possible to reduce ANN to approximate polytope membership [AFM11]
- Improved query time for storage between  $O(n/\epsilon^{d/4})$  and  $O(n/\epsilon^{d-1})$

# Bibliography

- [AFM11] S. Arya, G. D. da Fonseca, and D. M. Mount. Approximate polytope membership queries. *In Proc. 43rd Annu. ACM Sympos. Theory Comput.*, pages 579–586, 2011.
- [AMM09] S. Arya, T. Malamatos, and D. M. Mount. Space-time tradeoffs for approximate nearest neighbor searching. *J. ACM*, 57:1–54, 2009.
- [BI76] E. M. Bronshteyn and L. D. Ivanov. The approximation of convex sets by polyhedra. *Siberian Math. J.*, 16:852–853, 1976.
- [Dud74] R. M. Dudley. Metric entropy of some classes of sets with differentiable boundaries. *Approx. Theory*, 10(3):227–236, 1974.
- [Gru93] P. M. Gruber. Asymptotic estimates for best and stepwise approximation of convex bodies. *I. Forum Math.*, 5:521–537, 1993.
- [KR11] J. Kim and S. Reisner. Local minimality of the volume-product at the simplex. *Mathematika*, 57:121–134, 2011.
- [Kup08] G. Kuperberg. From the Mahler conjecture to Gauss linking integrals. *Geometric And Functional Analysis*, 18:870–892, 2008.

Thank you!