

Intersection Graphs of L-Shapes and Segments in the Plane^{*}

Stefan Felsner¹, Kolja Knauer², George B. Mertzios³, and Torsten Ueckerdt⁴

¹ Institut für Mathematik, Technische Universität Berlin, Germany.
Email: felsner@math.tu-berlin.de

² LIF UMR 7279, Université Aix-Marseille, CNRS, France.
Email: kolja.knauer@lif.univ-mrs.fr

³ School of Engineering and Computing Sciences, Durham University, UK.
Email: george.mertzios@durham.ac.uk

⁴ Department of Mathematics, Karlsruhe Institute of Technology, Germany.
Email: torsten.ueckerdt@kit.edu

Abstract. An L-shape is the union of a horizontal and a vertical segment with a common endpoint. These come in four rotations: L, Γ , J and \bar{J} . A k -bend path is a simple path in the plane, whose direction changes k times from horizontal to vertical. If a graph admits an intersection representation in which every vertex is represented by an L, an L or Γ , a k -bend path, or a segment, then this graph is called an $\{L\}$ -graph, $\{L, \Gamma\}$ -graph, B_k -VPG-graph or SEG-graph, respectively. Motivated by a theorem of Middendorf and Pfeiffer [Discrete Mathematics, 108(1):365–372, 1992], stating that every $\{L, \Gamma\}$ -graph is a SEG-graph, we investigate several known subclasses of SEG-graphs and show that they are $\{L\}$ -graphs, or B_k -VPG-graphs for some small constant k . We show that all planar 3-trees, all line graphs of planar graphs, and all full subdivisions of planar graphs are $\{L\}$ -graphs. Furthermore we show that complements of planar graphs are B_{17} -VPG-graphs and complements of full subdivisions are B_2 -VPG-graphs. Here a full subdivision is a graph in which each edge is subdivided at least once.

Keywords: intersection graphs, segment graphs, co-planar graphs, k -bend VPG-graphs, planar 3-trees.

1 Introduction and Motivation

A **segment intersection graph**, SEG-graph for short, is a graph that can be represented as follows. Vertices correspond to straight-line segments in the plane and two vertices are adjacent if and only if the corresponding segments intersect. Such representations are called *SEG-representations* and, for convenience, the class of all SEG-graphs is denoted by SEG. SEG-graphs are an important subject

^{*} This work was partially supported by (i) the DFG ESF EuroGIGA projects COMPOSE and GraDR, (ii) the EPSRC Grant EP/K022660/1, (iii) the ANR Project EGOS: ANR-12-JS02-002-01, and (iv) the PEPS grant EROS. A preliminary version appeared in the Proceedings of MFCS 2014 [13].

of study strongly motivated from an algorithmic point of view. Indeed, having an intersection representation of a graph (in applications graphs often come along with such a given representation) may allow for designing better or faster algorithms for optimization problems that are hard for general graphs, such as finding a maximum clique in interval graphs.

More than 20 years ago, Middendorf and Pfeiffer [24], considered intersection graphs of **axis-aligned L-shapes** in the plane, where an axis-aligned L-shape is the union of a horizontal and a vertical segment whose intersection is an endpoint of both. In particular, L-shapes come in four possible rotations: \mathbf{L} , $\mathbf{\Gamma}$, \mathbf{J} , and $\mathbf{7}$. For a subset X of these four rotations, e.g., $X = \{\mathbf{L}\}$ or $X = \{\mathbf{L}, \mathbf{\Gamma}\}$, we call a graph an X -graph if it admits an X -representation, i.e., vertices can be represented by L-shapes from X in the plane, each with a rotation from X , such that two vertices are adjacent if and only if the corresponding L-shapes intersect. Similarly to SEG, we denote the class of all X -graphs by X . The question if an intersection representation with polygonal paths or pseudo-segments can be *stretched* into a SEG-representation is a classical topic in combinatorial geometry and Oriented Matroid Theory. Middendorf and Pfeiffer prove the following interesting relation between intersection graphs of segments and L-shapes.

Theorem 1 (Middendorf and Pfeiffer [24]). *Every $\{\mathbf{L}, \mathbf{\Gamma}\}$ -representation has a combinatorially equivalent SEG-representation.*

This theorem is best-possible in the sense that there are examples of $\{\mathbf{L}, \mathbf{7}\}$ -graphs which are no SEG-graphs [6, 24], i.e., such $\{\mathbf{L}, \mathbf{7}\}$ -representations cannot be stretched. We feel that Theorem 1, which of course implies that $\{\mathbf{L}, \mathbf{\Gamma}\} \subseteq \text{SEG}$, did not receive a lot of attention in the active field of SEG-graphs. In particular, one could use Theorem 1 to prove that a certain graph class \mathcal{G} is contained in SEG by showing that \mathcal{G} is contained in $\{\mathbf{L}, \mathbf{\Gamma}\}$. For example, very recently Pawlik *et al.* [25] discovered a class of triangle-free SEG-graphs with arbitrarily high chromatic number, disproving a famous conjecture of Erdős [18], and it is in fact easier to see that these graphs are $\{\mathbf{L}\}$ -graphs than to see that they are SEG-graphs. To the best of our knowledge, the stronger result $\mathcal{G} \subseteq \{\mathbf{L}, \mathbf{\Gamma}\}$ has never been shown for any non-trivial graph class \mathcal{G} . In this paper we initiate this research direction. We consider several graph classes which are known to be contained in SEG and show that they are actually contained in $\{\mathbf{L}\}$, which is a proper subclass of $\{\mathbf{L}, \mathbf{\Gamma}\}$ [6].

Whenever a graph is not known (or known not) to be an intersection graph of segments or axis-aligned L-shapes, one often considers natural generalizations of these intersection representations. Asinowski *et al.* [3] introduced **intersection graphs of axis-aligned k -bend paths** in the plane, called B_k -VPG-graphs. An (axis-aligned) k -bend path is a simple path in the plane, whose direction changes k times from horizontal to vertical. Clearly, B_1 -VPG-graphs are precisely intersection graphs of all four L-shapes; the union of B_k -VPG-graphs for all $k \geq 0$ is exactly the class STRING of intersection graphs of simple curves in the plane [3]. Now if a graph $G \notin \text{SEG}$ is a B_k -VPG-graph for some small k , then one might say that G is “not far from being a SEG-graph”.

Our Results and Related Work.

Let us denote the class of all planar graphs by PLANAR. A recent celebrated result of Chalopin and Gonçalves [5] states that $\text{PLANAR} \subset \text{SEG}$, which was conjectured by Scheinerman [26] in 1984. However, their proof is rather involved and there is not much control over the kind of SEG-representations. Here we give an easy proof for a non-trivial subclass of planar graphs, namely *planar 3-trees*. A *3-tree* is an edge-maximal graph of treewidth 3. Every 3-tree can be built up starting from the clique K_4 and adding new vertices, one at a time, whose neighborhood in the so-far constructed graph is a triangle.

Theorem 2. *Every planar 3-tree is an $\{L\}$ -graph.*

It remains open to generalize Theorem 2 to planar graphs of treewidth 3 (i.e., subgraphs of planar 3-trees). On the other hand it is easy to see that graphs of treewidth at most 2 are $\{L\}$ -graphs [7]. Chaplick and the last author show in [8] that planar graphs are B_2 -VPG-graphs, improving on an earlier result of Asinowski *et al.* [3]. In [8] it is also conjectured that $\text{PLANAR} \subset \{L\}$, which with Theorem 1 would imply the main result of [5], i.e., $\text{PLANAR} \subset \text{SEG}$.

Considering line graphs of planar graphs, one easily sees that these graphs are SEG-graphs. Indeed, a straight-line drawing of a planar graph G can be interpreted as a SEG-representation of the line graph $L(G)$ of G , which has the edges of G as its vertices and pairs of incident edges as its edges. We prove the following strengthening result.

Theorem 3. *The line graph of every planar graph is an $\{L\}$ -graph.*

Kratochvíl and Kuběna [21] consider the class of complements of planar graphs (co-planar graphs), CO-PLANAR for short. They show that every graph in CO-PLANAR is an intersection graph of convex sets in the plane, and ask whether $\text{CO-PLANAR} \subset \text{SEG}$. As the INDEPENDENT SET PROBLEM in planar graphs is known to be NP-complete [15], MAX CLIQUE is NP-complete for any graph class $\mathcal{G} \supseteq \text{CO-PLANAR}$, e.g., intersection graphs of convex sets. Indeed, the longstanding open question whether MAX CLIQUE is NP-complete for SEG [22] has recently been answered affirmatively by Cabello, Cardinal and Langerman [4] by showing that every planar graph has an even subdivision whose complement is a SEG-graph. The subdivision is essential in the proof of [4], as it still remains an open problem whether $\text{CO-PLANAR} \subset \text{SEG}$ [21]. The largest subclass of CO-PLANAR known to be in SEG is the class of complements of partial 2-trees [14]. Here we show that all co-planar graphs are “not far from being SEG-graphs”.

Theorem 4. *Every co-planar graph is a B_{17} -VPG-graph.*

Theorem 4 implies that MAX CLIQUE is NP-complete for B_k -VPG-graphs with $k \geq 17$. On the other hand, the MAX CLIQUE problem for B_0 -VPG-graphs can be solved in polynomial time, while VERTEX COLORABILITY remains NP-complete but allows for a 2-approximation [3]. Middendorf and Pfeiffer [24] show that the complement of any *even subdivision* of any graph, i.e., every edge is

subdivided with a non-zero even number of vertices, is an $\{L, \Gamma\}$ -graph. This implies that MAX CLIQUE is NP-complete even for $\{L, \Gamma\}$ -graphs.

We consider *full subdivisions* of graphs, that is, a subdivision H of a graph G where each edge of G is subdivided at least once. It is not hard to see that a full subdivision H of G is in STRING if and only if G is planar, and that if G is planar, then H is actually a SEG-graph. Here we show that this can be further strengthened, namely that H is in an $\{L\}$ -graph. Moreover, we consider the complement of a full subdivision H of an arbitrary graph G , which is in STRING but not necessarily in SEG. Here, similar to the result of Middendorf and Pfeiffer [24] on even subdivisions we show that such a graph H is “not far from being SEG-graph”.

Theorem 5. *Let H be a full subdivision of a graph G .*

- (i) *If G is planar, then H is an $\{L\}$ -graph.*
- (ii) *If G is any graph, then the complement of H is a B_2 -VPG-graph.*

The graph classes considered in this paper are illustrated in Figure 1. We shall prove Theorems 2, 3, 4 and 5 in Sections 2, 3, 4 and 5, respectively, and conclude with some open questions in Section 6.




Fig. 1. Graph classes considered in this paper.

Related Representations.

In the context of *contact representations*, where distinct segments or k -bend paths may not share interior points, it is known that every contact SEG-representation

has a combinatorially equivalent contact B_1 -VPG-representation, but not vice versa [20]. Contact SEG-graphs are exactly planar Laman graphs and their subgraphs [10], which includes for example all triangle-free planar graphs. Very recently, contact $\{L\}$ -graphs have been characterized [7]. Necessary and sufficient conditions for stretchability of a contact system of pseudo-segments are known [1, 11].

Let us also mention the closely related concept of *edge*-intersection graphs of paths in a grid (EPG-graphs) introduced by Golumbic *et al.* [16]. There are some notable differences, starting from the fact that *every* graph is an EPG-graph [16]. Nevertheless, analogous questions to the ones posed about VPG-representations of STRING-graphs are posed about EPG-representations of general graphs. In particular, there is a strong interest in finding representations using paths with few bends, see [19] for a recent account.

2 Proof of Theorem 2

Proof. Let G be a plane 3-tree with a fixed plane embedding. We construct an $\{L\}$ -representation of G satisfying the additional property that for every inner triangular face $\{a, b, c\}$ of G there exists a subset of the plane, called the *private region of the face*, that intersects only the L -paths for a , b and c , and no other L -path. More precisely, a private region of $\{a, b, c\}$ is an axis-aligned rectilinear polygon having one of the shapes depicted in Figure 2(a), such that the L -paths for a , b and c intersect the polygon as shown in figure.




Fig. 2. (a) The two possible shapes of a private region for inner facial triangle $\{a, b, c\}$. (b) An $\{L\}$ -representation of the plane 3-tree on three vertices together with a private region for the only inner face.

Indeed, we prove the following stronger statement by induction on the number of vertices in G .

Claim. Every plane 3-tree admits an $\{L\}$ -representation together with a private region for every inner face, such that the private regions for distinct faces are disjoint.

As induction base ($|V(G)| = 3$) consider the graph G consisting only of the triangle $\{a, b, c\}$. Then there is an essentially unique $\{L\}$ -representation of G and it is not difficult to find a private region for the unique inner face of G . We refer to Figure 2(b) for an illustration.

Now let us assume that $|V(G)| \geq 4$. Because G is a 3-tree there exists an inner vertex v of degree exactly three. In particular, the three neighbors a, b, c of v form an inner facial triangle in the plane 3-tree $G' = G \setminus v$. By induction G' admits an $\{L\}$ -representation with a private region for each inner face so that distinct private regions are disjoint.

Consider the private region R for $\{a, b, c\}$. By flipping the plane along the main diagonal if necessary, we can assume without loss of generality that R has the shape shown in the left of Figure 2(a). (Note that such a flip does not change the type of the L-paths.) Now we introduce an L-path for vertex v completely inside R as depicted in Figure 3(a). Since R does not intersect any other L-path this is an $\{L\}$ -representation of G .




Fig. 3. (a) Introducing an L-shape for vertex v into the private region for the triangle $\{a, b, c\}$. (b) Identifying a pairwise disjoint private regions for the facial triangles $\{a, b, v\}$, $\{a, c, v\}$ and $\{b, c, v\}$.

Finally we identify three private regions for the three newly created inner faces $\{a, b, v\}$, $\{a, c, v\}$ and $\{b, c, v\}$. This is shown in Figure 3(b). Since these regions are pairwise disjoint and completely contained in the private region for $\{a, b, c\}$ we have identified a private region for every inner face so that distinct regions are disjoint. (Note that $\{a, b, c\}$ is not a facial triangle in G and hence does not need a private region.) This proves the claim and thus concludes the proof of the theorem. \square

3 Proof of Theorem 3

Proof. Without loss of generality let G be a maximally planar graph with a fixed plane embedding. (Line graphs of subgraphs of G are induced subgraphs of $L(G)$.) Then G admits a so-called *canonical ordering* –first defined in [12]–, namely an ordering v_1, \dots, v_n of the vertices of G such that

- Vertices v_1, v_2, v_n form the outer triangle of G in clockwise order. (We draw G such that v_1, v_2 are the highest vertices.)

- For $i = 3, \dots, n$ vertex v_i lies in the outer face of the induced embedded subgraph $G_{i-1} = G[v_1, \dots, v_{i-1}]$. Moreover, the neighbors of v_i in G_{i-1} form a path on the outer face of G_{i-1} with at least two vertices.

We shall construct an $\{\mathbf{L}\}$ -representation of $L(G)$ along a fixed canonical ordering v_1, \dots, v_n of G . For every $i = 2, \dots, n$ we shall construct an $\{\mathbf{L}\}$ -representation of $L(G_i)$ with the following additional properties.

For every outer vertex v of G_i we maintain an auxiliary bottomless rectangle $R(v)$, i.e., an axis-aligned rectangle with bottom-edge at $-\infty$, such that:

- $R(v)$ intersects the horizontal segments of precisely those rectilinear paths for edges in G_i incident to v .
- $R(v)$ does not contain any bends or endpoints of any path for an edge in G_i and does not intersect any $R(w)$ for $w \neq v$.
- the left-to-right order of the bottomless rectangles matches the order of vertices on the counterclockwise outer (v_1, v_2) -path of G_i .

The bottomless rectangles act as placeholders for the upcoming vertices of $L(G)$. Indeed, all upcoming intersections of paths will be realized inside the corresponding bottomless rectangles. For $i = 2$, the graph G_i consist only of the edge $\{v_1, v_2\}$. Hence an $\{\mathbf{L}\}$ -representation of the one-vertex graph $L(G_2)$ consists of only one L-shape and two disjoint bottomless rectangles $R(v_1)$, $R(v_2)$ intersecting its horizontal segment.

For $i \geq 3$, we start with an $\{\mathbf{L}\}$ -representation of $L(G_{i-1})$. Let (w_1, \dots, w_k) be the counterclockwise outer path of G_{i-1} that corresponds to the neighbors of v_i in G_{i-1} . The corresponding bottomless rectangles $R(w_1), \dots, R(w_k)$ appear in this left-to-right order. See Figure 4 for an illustration. For every edge $\{v_i, w_j\}$, $j = 1, \dots, k$ we define an L-shape $P(v_i w_j)$ whose vertical segment is contained in the interior of $R(w_j)$ and whose horizontal segment ends in the interior of $R(w_k)$. Moreover, the upper end and lower end of the vertical segment of $P(v_i w_j)$ lies on the top side of $R(w_j)$ and below all L-shapes for edges in G_{i-1} , respectively. Finally, the bend and right end of $P(v_i w_j)$ is placed above the bend of $P(v_i w_{j+1})$ and to the right of the right end of $P(v_i w_{j+1})$ for $j = 1, \dots, k - 1$, see Figure 4.

It is straightforward to check that this way we obtain an $\{\mathbf{L}\}$ -representation of $L(G_i)$. So it remains to find a set of bottomless rectangles, one for each outer vertex of G_i , satisfying our additional property. We set $R'(v) = R(v)$ for every $v \in V(G_i) \setminus \{v_i, w_1, \dots, w_k\}$ since these are kept unchanged. Since $R(w_1)$ and $R(w_k)$ are not valid anymore, we define a new bottomless rectangle $R'(w_1) \subset R(w_1)$ such that $R'(w_1)$ is crossed by all horizontal segments that cross $R(w_1)$ and additionally the horizontal segment of $P(v_i w_1)$. Similarly, we define $R'(w_k) \subset R(w_k)$. And finally, we define a new bottomless rectangle $R'(v_i) \subset R(w_k)$ in such a way that it is crossed by the horizontal segments of exactly $P(v_i w_1), \dots, P(v_i w_k)$. Note that for $1 < j < k$ the outer vertex w_j of G_{i-1} is not an outer vertex of G_i . Then $\{R'(v) \mid v \in v(G_i)\}$ has the desired property. See again Figure 4. \square




Fig. 4. Along a canonical ordering a vertex v_i is added to G_{i-1} . For each edge between v_i and a vertex in G_{i-1} an L-shape is introduced with its vertical segment in the corresponding bottomless rectangle. The three new bottomless rectangles $R'(w_1), R'(v_i), R'(w_4)$ are highlighted.

4 Proof of Theorem 4

Proof. Let $G = (V, E)$ be any planar graph. We shall construct a B_k -VPG-representation of the complement \bar{G} of G for some constant k that is independent of G . Indeed, $k = 17$ is enough. To find the VPG-representation we make use of two crucial properties of G : A) G is 4-colorable and B) G is 5-degenerate. Indeed, our construction gives a B_{2d+7} -VPG-representation for the complement of any 4-colorable d -degenerate graph. Here a graph is called *d-degenerate* if it admits a vertex ordering such that every vertex has at most d neighbors with smaller index.

Consider any 4-coloring of G with color classes V_1, V_2, V_3, V_4 . Further let v_1, \dots, v_n be an order of the vertices of V witnessing the degeneracy of G , i.e., for each v_i there are at most 5 neighbors v_j of v_i with $j < i$. We call these neighbors the *back neighbors* of v_i . Further consider the axis-aligned rectangle $R = [0, 2(n+1)] \times [0, n+1]$.

Consider any ordered pair of color classes, say (V_1, V_2) and the axis-aligned rectangle $R = [0, 2(n+1)] \times [0, n+1]$. We will represent all non-edges of the form (v_i, v_j) with $i < j$, $v_i \in V_1$, and $v_j \in V_2$ in R . We define a monotone increasing path $Q(v)$ for each $v \in V_1 \cup V_2$ as follows. See Figure 5 for an illustration.

- For $v_j \in V_1$ let $Q(v_j)$ start on $(0, j)$, go horizontally to $(n+1+j, j)$ and then vertically to $(n+1+j, n+1+j)$. (These are the black paths in the middle of Figure 5.)
- If $v_j \in V_2$ with back neighbors $v_{i_1}, \dots, v_{i_{k'}} \in V_1$ let $Q(v_j)$ start vertically on $(j, 0)$, bend to the right at $(j, i_1 - \varepsilon)$, bend upwards at $(n+1+i_1 + \varepsilon, i_1 - \varepsilon)$, and in this way avoid all paths representing back neighbors of v_j in V_1 . More precisely, $Q(v_j)$ now uses horizontal segments of the form $[(n+1+i_\ell + \varepsilon, i_{\ell+1} - \varepsilon), (n+1+i_{\ell+1} + \varepsilon, i_{\ell+1} - \varepsilon)]$ and vertical segments of the form $[(n+1+i_\ell + \varepsilon, i_\ell - \varepsilon), (n+1+i_\ell + \varepsilon, i_{\ell+1} - \varepsilon)]$, for $\ell + 1 \leq k'$. The last vertical segment of $Q(v_j)$ starts at $[(n+1+i_{k'} + \varepsilon, i_{k'} - \varepsilon)]$ and goes




Fig. 5. Left: The induced subgraph $G[V_1, V_2]$ for a pair of color classes (V_1 (black), V_2 (white)) of a planar graph G . Middle: A VPG-representation of all non-edges of the form (v_i, v_j) with $i < j$, $v_i \in V_1$, and $v_j \in V_2$ in R . Right: A schematic representation of the arrangement of paths. T arrows indicated the order of coordinates within V_1 and V_2 . The disk on the bend of the black path indicates, that backward non-edges to V_1 are realized.

to $(n + 1 + i_\ell + \varepsilon, j)$. From there $Q(v_j)$ takes the last horizontal segment to $(2(n + 1), j)$. (These are the gray paths in the middle of Figure 5.)

First, note that in order to avoid collinearities for each j there should be a different $\varepsilon(j)$ in the above construction, which yields a representation on an integer grid of size roughly $2n^2 \times n^2$.

Second, note that all paths representing vertices from V_1 mutually cross using one bend each, i.e., non-edges between vertices of V_1 are represented. Now observe that $\{Q(v) \mid v \in V_1 \cup V_2\}$ is a VPG-representation of all non-edges of the form (v_i, v_j) with $i < j$, $v_i \in V_1$, and $v_j \in V_2$. Every path $Q(v_j)$ with $v_j \in V_2$ crosses all paths corresponding to $v_i \in V_1$ with $i < j$ except those that are back neighbors of v_j . These are avoided at a cost of two bends per back neighbor. A last bend is spent when turning to the right the last time.

Furthermore, note that the order on the y -coordinates of the starting points of the V_1 -paths and the order on the x -coordinates of the last points of the V_1 -paths are according to their order of indices. Also observe that the order of the x -coordinates of the starting points of the V_2 -paths does not matter for the construction, as long as all of them are in $[0, n + 1]$. Last, note that the order on the y -coordinates of the last points of the V_2 -paths is according to their order of indices. We synthesize this situation into a diagram on the right of Figure 5, where paths represent the family of paths from one color class, the arrow indicates the order in which the paths within one family are arranged. The disk on the bend of the black path means, that the non-edges from the gray class backwards to the black class are realized.

Now we have defined for each ordered pair of color classes (V_i, V_j) a VPG-representation of the non-edges backward from V_j to V_i . For every vertex $v \in V$ we have defined six Q -paths, two for each colors class that v is not in. In total the six Q -paths for the same vertex v have at most $6 + 2k \leq 16$ bends, where $k \leq 5$ is the back degree of v . It remains to place (translate and rotate) the

six representations into non-overlapping positions and to “connect” the three Q -paths for each vertex in such a way that connections for vertices of different color do not intersect. This can be done with at most three extra bends per path, see Figure 6. Finally, note that the first and last segment of every path in the representation can be omitted, yielding the claimed bound. \square




Fig. 6. Interconnecting the VPG-representations by adding at most three bends for each vertex. The set of paths corresponding to color class V_i is indicated by a single path labeled V_i , $i = 1, 2, 3, 4$. Little gray boxes on bends indicate that the bend is used to reverse the order of coordinates within the class.

5 Proof of Theorem 5

Proof. Let G be any graph and H arise from G by subdividing each edge at least once. Without loss of generality we may assume that every edge of G is subdivided exactly once or twice. Indeed, if an edge e of G is subdivided three times or more, then H can be seen as a full subdivision of the graph G' that arises from G by subdividing e once.

- (i) Assuming that G is planar, we shall find an $\{L\}$ -representation of H as follows. Without loss of generality G is maximally planar. We consider a bar visibility representation of G , i.e., vertices of G are disjoint horizontal segments in the plane and edges are disjoint vertical segments in the plane whose endpoints are contained in the two corresponding vertex segments and which are disjoint from all other vertex segments. Such a representation for a planar triangulation exists e.g. by [23]. See Figure 7 for an illustration.




Fig. 7. A planar graph G on the left, a bar visibility representation of G in the center, and an $\{L\}$ -representation of a full division of G on the right. Here, the edges $\{1, 2\}$, $\{1, 3\}$ and $\{3, 6\}$ are subdivided twice.

It is now easy to interpret every segment as an L, and replace a segment corresponding to an edge that is subdivided twice by two L-shapes. Let us simply refer to Figure 7 again.

- (ii) Now assume that $G = (V, E)$ is any graph. We shall construct a B_2 -VPG-representation of the complement \bar{H} of $H = (V \cup W, E')$ with monotone increasing paths only. First, we represent the clique $\bar{H}[V]$. Let $V = \{v_1, \dots, v_n\}$ and define for $i = 1, \dots, n$ the 2-bend path $P(v_i)$ for vertex v_i to start at $(i, 0)$, have bends at (i, i) and $(i+n, i)$, and end at $(i+n, n+1)$. See Figure 8 for an illustration. For convenience, let us call these paths v -paths.

Next, we define for every edge of G the 2-bend paths for the one or two corresponding subdivision vertices in \bar{H} . We call these paths w -paths. So let $\{v_i, v_j\}$ be any edge of G with $i < j$. We distinguish two cases.

- Case 1.* The edge $\{v_i, v_j\}$ is subdivided by only one vertex w_{ij} in H . We define the w -path $P(w_{ij})$ to start at $(j - \frac{1}{4}, i + \frac{1}{4})$, have bends at $(j - \frac{1}{4}, j + \frac{1}{4})$ and $(i+n - \frac{1}{4}, j + \frac{1}{4})$, and end at $(i+n - \frac{1}{4}, n+1)$, see the left of Figure 8.
- Case 2.* The edge $\{v_i, v_j\}$ is subdivided by two vertices w_i, w_j with $\{v_i, w_i\}, \{v_j, w_j\} \in E(H)$. We define the start, bends and end of the w -path $P(w_i)$ to be $(j - \frac{1}{4}, i + \frac{1}{4})$, $(j - \frac{1}{4}, j - \frac{1}{4})$, $(i+n - \frac{1}{4}, j - \frac{1}{4})$ and $(i+n - \frac{1}{4}, n+1)$, respectively. The start, bends and end of the w -path $P(w_j)$ are $(j - \frac{1}{2}, i - \frac{1}{4})$, $(j - \frac{1}{2}, j + \frac{1}{4})$, $(i+n - \frac{1}{2}, j + \frac{1}{4})$ and $(i+n - \frac{1}{2}, n+1)$, respectively. See the right of Figure 8.

It is easy to see that every w -path $P(w)$ intersects every v -path, except for the one or two v -paths corresponding to the neighbors of w in H . Moreover,




Fig. 8. Left: Inserting the path $P(w_{ij})$ for a single vertex w_{ij} subdividing the edge $v_i v_j$ in G . Right: Inserting the paths $P(w_i)$ and $P(w_j)$ for two vertices w_i, w_j subdividing the edge $v_i v_j$ in G .

the two w -paths in Case 2 are disjoint. It remains to check that the w -paths for distinct edges of G mutually intersect. To this end, note that every w -path for edge $v_i v_j$ starts near (j, i) , bends near (j, j) and $(i + n, j)$ and ends near $(i + n, n)$. Consider two w -paths P and P' that start at (j, i) and (j', i') , respectively, and bend near (j, j) and (j', j') , respectively. If $j = j'$ then it is easy to check that P and P' intersect near (j, j) . Otherwise, let $j' > j$. Now if $j > i'$, then P and P' intersect near (j', i) , and if $j \leq i'$, then P and P' intersect near $(i + n, j')$.

Hence we have found a B_2 -VPG-representation of \bar{H} , as desired. Let us remark, that in this representation some w -paths intersect non-trivially along some horizontal or vertical lines, i.e., share more than a finite set of points. However, this can be omitted by a slight and appropriate perturbation of endpoints and bends of w -paths. \square

6 Conclusions and Open Problems

Motivated by Middendorf and Pfeiffer's theorem (Theorem 1 in [24]) that every $\{\mathbb{L}, \Gamma\}$ -representation can be stretched into a SEG-representation, we considered the question which subclasses of SEG-graphs are actually $\{\mathbb{L}, \Gamma\}$ -graphs, or even $\{\mathbb{L}\}$ -graphs. We proved that this is indeed the case for several graph classes related to planar graphs. We feel that the question whether $\text{PLANAR} \subset \{\mathbb{L}, \Gamma\}$, as already conjectured [8], is of particular importance. After all, this, together with Theorem 1, would give a new proof for the fact that $\text{PLANAR} \subset \text{SEG}$.

Open Problem. *Each of the following is open.*

- (i) *When can a B_1 -VPG-representation be stretched into a combinatorially equivalent SEG-representation?*
- (ii) *Is $\{\mathbb{L}, \Gamma\} = \text{SEG} \cap B_1\text{-VPG}$?*

- (iii) Is every planar graph an $\{\mathbb{L}\}$ -graph, or B_1 -VPG-graph?
- (iv) Does every planar graph admit an even subdivision whose complement is an $\{\mathbb{L}\}$ -graph, or B_1 -VPG-graph?
- (v) Recognizing B_k -VPG-graphs is known to be NP-complete for each $k \geq 0$ [6]. What is the complexity of recognizing $\{\mathbb{L}\}$ -graphs, or $\{\mathbb{L}, \Gamma\}$ -graphs?

Recall that $\cup_{k \geq 0} B_k\text{-VPG} = \text{STRING}$ [3]. Chaplick *et al.* [6] prove that $B_k\text{-VPG} \subsetneq B_{k+1}\text{-VPG}$ for all $k \geq 0$ and also that $\text{SEG} \not\subseteq B_k\text{-VPG}$ for each $k \geq 0$, even if SEG is restricted to three slopes only. Another natural subclass of STRING, which is in no inclusion-relation with SEG, is the class COCOMP of co-comparability graphs [17]. However, one can prove a result similar to the previous one concerning B_k -VPG-graphs and STRING-graphs:

There is no $k \in \mathbb{N}$ such that $B_k\text{-VPG} \supset \text{COCOMP}$, compare also Figure 1. A proof can be given along the “degrees of freedoms” approach of Alon and Scheinerman [2], i.e., by counting the graphs in the respective sets.

First, Alon and Scheinerman consider the class $P(n, t)$ of t -dimensional posets on n elements and show that for fixed t the growth of $\log |P(n, t)|$ behaves like $nt \log n$. If $CC(n, t)$ denotes the class of cocomparability graphs of posets in $P(n, t)$, then with easy adaptations we obtain $\log |CC(n, t)| \geq n(t - 1 - o(1)) \log n$.

On the other hand, every path of a B_k -VPG-representation can be encoded by $k + 4$ numbers. Whether two paths intersect can be answered by looking at the signs of few low degree polynomials in $2k + 8$ variables evaluated at the encodings of the two paths, meaning that the class B_k -VPG has $k + 4$ degrees of freedom. Alon and Scheinerman show how to use Warren’s Theorem [27] to get an upper bound on the size of such a class. Indeed, the logarithm of the number of B_k -VPG-graphs on n vertices is at most $O(1)nk \log n$.

Comparing the numbers we find that there is a cocomparability graph of a $(k + 2)$ -dimensional poset that is not a B_k -VPG-graph. On the other hand it is easy to find a B_k -VPG-representation for the cocomparability graph of any given $(k + 1)$ -dimensional poset. So we have for every $k \in \mathbb{N}$

$$\bigcup_{n \in \mathbb{N}} CC(n, k + 1) \subseteq B_k\text{-VPG} \not\supseteq \bigcup_{n \in \mathbb{N}} CC(n, k + 2).$$

Recently, this has been proved independently by Cohen, Golumbic, Trotter and Wang [9] and hence we omit more details here.

Comparing this to the result that there is no k such that $\text{SEG} \subseteq B_k\text{-VPG}$ a natural question arises. Is there a parameter of SEG-graphs or SEG-representations ensuring few bends in their VPG-representations? It is known that the number of slopes in the SEG-representation is not the right answer [6].

References

1. N. Aerts and S. Felsner. Straight line triangle representations. In *Proceedings of the 21st International Symposium on Graph Drawing (GD)*, pages 119–130, 2013.

2. N. Alon and E. Scheinerman. Degrees of freedom versus dimension for containment orders. *Order*, 5:11–16, 1988.
3. A. Asinowski, E. Cohen, M. C. Golumbic, V. Limouzy, M. Lipshteyn, and M. Stern. Vertex intersection graphs of paths on a grid. *J. Graph Algorithms Appl.*, 16(2):129–150, 2012.
4. S. Cabello, J. Cardinal, and S. Langerman. The clique problem in ray intersection graphs. *Discrete & Computational Geometry*, 50(3):771–783, 2013.
5. J. Chalopin and D. Gonçalves. Every planar graph is the intersection graph of segments in the plane: extended abstract. In *Proceedings of the 41st annual ACM symposium on Theory of computing*, STOC '09, pages 631–638, 2009.
6. S. Chaplick, V. Jelínek, J. Kratochvíl, and T. Vyskočil. Bend-bounded path intersection graphs: Sausages, noodles, and waffles on a grill. In *Proceedings of the 38th Graph-Theoretic Concepts in Computer Science (WG)*, pages 274–285, 2012.
7. S. Chaplick, S. G. Kobourov, and T. Ueckerdt. Equilateral L-contact graphs. In *Proceedings of the 39th Graph-Theoretic Concepts in Computer Science (WG)*, pages 139–151, 2013.
8. S. Chaplick and T. Ueckerdt. Planar graphs as VPG-graphs. *Journal of Graph Algorithms and Applications*, 17(4):475–494, 2013.
9. E. Cohen, M. C. Golumbic, W. T. Trotter, and R. Wang. Posets and VPG graphs. <http://people.math.gatech.edu/~trotter/papers/140.pdf>, 2014.
10. H. de Fraysseix and P. O. de Mendez. Representations by contact and intersection of segments. *Algorithmica*, 47(4):453–463, 2007.
11. H. de Fraysseix and P. O. de Mendez. Stretching of Jordan arc contact systems. *Discrete Applied Mathematics*, 155(9):1079–1095, 2007.
12. H. De Fraysseix, J. Pach, and R. Pollack. How to draw a planar graph on a grid. *Combinatorica*, 10(1):41–51, 1990.
13. S. Felsner, K. Knauer, G. B. Mertzios, and T. Ueckerdt. Intersection graphs of L-shapes and segments in the plane. In *Proceedings of the 39th annual symposium on Mathematical Foundations of Computer Science*, MFCS '14, 2014.
14. M. C. Francis, J. Kratochvíl, and T. Vyskočil. Segment representation of a subclass of co-planar graphs. *Discrete Mathematics*, 312(10):1815–1818, 2012.
15. M. R. Garey and D. S. Johnson. *Computers and Intractability: A guide to the theory of NP-completeness*. W. H. Freeman, 1979.
16. M. C. Golumbic, M. Lipshteyn, and M. Stern. Edge intersection graphs of single bend paths on a grid. *Networks*, 54(3):130–138, 2009.
17. M. C. Golumbic, D. Rotem, and J. Urrutia. Comparability graphs and intersection graphs. *Discrete Mathematics*, 43(1):37–46, 1983.
18. A. Gyárfás. Problems from the world surrounding perfect graphs. *Zastos. Mat.*, 19(3-4):413–441, 1987.
19. D. Heldt, K. Knauer, and T. Ueckerdt. Edge-intersection graphs of grid paths: The bend-number. *Discrete Appl. Math.*, 167:144–162, 2014.
20. S. G. Kobourov, T. Ueckerdt, and K. Verbeek. Combinatorial and geometric properties of planar Laman graphs. In *SODA*, pages 1668–1678. SIAM, 2013.
21. J. Kratochvíl and A. Kuběna. On intersection representations of co-planar graphs. *Discrete Math.*, 178(1-3):251–255, 1998.
22. J. Kratochvíl and J. Matousek. Intersection graphs of segments. *Journal of Combinatorial Theory, Series B*, 62(2):289–315, 1994.
23. F. Luccio, S. Mazzone, and C. K. Wong. A note on visibility graphs. *Discrete Math.*, 64(2-3):209–219, 1987.
24. M. Middendorf and F. Pfeiffer. The max clique problem in classes of string-graphs. *Discrete mathematics*, 108(1):365–372, 1992.

25. A. Pawlik, J. Kozik, T. Krawczyk, M. Lasoń, P. Micek, W. T. Trotter, and B. Walczak. Triangle-free intersection graphs of line segments with large chromatic number. *Journal of Combinatorial Theory, Series B*, 2013.
26. E. R. Scheinerman. *Intersection classes and multiple intersection parameters of graphs*. PhD thesis, Princeton University, 1984.
27. H. E. Warren. Lower bounds for approximation by nonlinear manifolds. *Trans. Amer. Math. Soc.*, 133:167–178, 1968.